

Appendix 1

Industrial Site Photos: taken from 2016 to 2020

2012 Herbicide Application:

Observed site composition in 2018 consisted of mullein, St. John's wort, Canadian thistle, spotted knapweed, willow, wood rose, and Oregon grape.



2017



2018



2020

2014 (1) Herbicide Treatment:

Observed site composition in 2018 consisted of Canadian thistle, spotted knapweed, and spirea.



2016



2017



2018

2014 (2) Herbicide Treatment:

Observed site composition in summer 2020 consisted of spotted knapweed, Canadian thistle, oxeyed daisy, and willow.



2017



2018



2019



2020

2014 (3) Herbicide treatment:

Observed site composition in summer 2020 consists of oxeye daisy, St. John's wort, golden rod, willow, and Rocky Mountain maple. Site was planted in 2017.



2016 Summer



2017 Summer



2019 Summer



2020 Summer

2015 (1) Herbicide Treatment:

Observed site composition in summer 2020 consists of knapweed, Canadian thistle, mullein, spirea, and willow.



2016



2018



2020

2015 (2) herbicide application:

Observed site composition in summer 2020 consists of spotted knapweed, Canadian thistle, mullein, St. John's wort and willow. Planted 2018



2016



2017



2019



2020

2016 (1) herbicide application:

Observed site composition in summer 2020 consists of knapweed, mullein, and Canadian horseweed.



2017



2018



2019



2020

State and Federal Lands Adjacent to 2016 (1):

Composition consisting of fireweed, ocean spray, and red stem ceanothus



Summer 2020



Summer 2020

2016 (2) and 2020 Herbicide Application:

Observed site composition in summer 2020 consists of mullein, prickly lettuce, and spotted knapweed.



2016



2017



2018



2018

2016 (3) and 2020 Herbicide Applications:

Observed site composition in summer 2020 consists of knapweed, Canadian thistle, and prickly lettuce.



2016



2017



2018



Spring 2020



Summer 2020

2016 (4) Herbicide Application:

Observed site composition in summer 2020 consists of knapweed, Canadian thistle, mullein, fireweed, Canadian horseweed.



2017



2018



2019



2020

2016 (5) Herbicide Application:

Observed site composition in summer 2020 consists of knapweed, Canadian thistle, mullein, fireweed, Canadian horseweed.



2017



2019



2020

2016 (6) Herbicide Application Post 2015 Fire:

Observed site composition in summer 2020 consists of knapweed, St. John's wort, mullein, and fireweed.



2016



2019



2020

Land adjacent to 2016 () recovering from 2015 fire with shrub and trees component <10ft in height

